

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 235/18, A61K 31/415, C07D 401/12, A61K 31/445, C07D 471/04, 473/00

(11) International Publication Number:

WO 98/06703

(43) International Publication Date:

19 February 1998 (19.02.98)

(21) International Application Number:

PCT/US97/13870

A1

(22) International Filing Date:

6 August 1997 (06.08.97)

(30) Priority Data:

60/023,942

14 August 1996 (14.08.96) US

(71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): CONNOR, David, Thomas [GB/US]; 2453 Antietam, Ann Arbor, MI 48105 (US). GLASE, Shelly, Ann [US/US]; 4468 Hillside Court, Ann Arbor, MI 48105 (US). PURCHASE, Terri, Stoeber [US/US]; 4961 Ravine Court, Ann Arbor, MI 48105 (US). ROTH, Bruce, David [US/US]; 49255 Hunt Club Court, Plymouth, MI 48170 (US). TRIVEDI, Bharat, Kalidas [IN/US]; 36955 Aldgate Court, Farmington Hills, MI 48335 (US).
- (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.

(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, HU, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: 2-PHENYL BENZIMIDAZOLE DERIVATIVES AS MCP-1 ANTAGONISTS

(57) Abstract

Benzimidazole derivatives of formula (I) or a pharmaceutically acceptable salt thereof are MCP-1 antagonists and are thus useful in the treatment of inflammation, artherosclerosis, restenosis, and immune disorders such as arthritis and transplant rejection wherein A is N or CH; where W, X, Y and Z can be independently C-R2, C-R3, C-R4, C-R5, or N; no more than two of W, X, Y and Z can be N in any one structure, R2, R3, R4 and R5 can be independently H, C₁₋₂₀ alkyl, halogen, nitro, -SO₂NR₈R9, alkoxy of from 1-4 carbon atoms; -S(O)_pR where p is an integer from 0 to 2; -(CH₂)_mOR, -(CH₂)_mCOOR, -(CH₂)_mCOOR₈R9, -(CH₂)_mCOOR₈R9, -(CH₂)_mCOOR, or -CF₃; m is an integer of from 0 to 4, R is hydrogen, lower alkyl of from 1-4 carbon atoms, aryl of from 6-10 carbon atoms, or benzyl; R₁ can be H, lower alkyl of from 1-4 carbon atoms, or -(CH₂)_mPh; R₆ is alkyl of from 1-6 carbon atoms or R₇; R₇ is (CH₂)_nNR₁₀R₁₁; n is an integer from 2 to 6; R₈ and R₉ can be independently hydrogen, lower alkyl of from 1-4 carbon atoms, or can be taken together to form a ring of from 3-8 atoms having up to one additional heteroatom as O, S, SO₂, or N-R₁₂; R₁₀ and R₁₁ can independently be lower alkyl, -(CH₂)_mPh, unsubstituted or substituted with up to three R₂ substituents, or R₁₀ and R₁₁ can be taken together to form a ring of from 3-8 atoms which may contain oxygen or NR₁₂; R₁₂ is hydrogen, lower alkyl, -(CH₂)₁Ph, where Ph is phenyl unsubstituted or substituted with up to three R₂ substituents; t is an integer of from 0 to 2.